



# gelclair<sup>®</sup>

concentrated oral gel

## Directions for Use

Initiate treatment with Gelclair<sup>®</sup> as soon as signs and symptoms of mucositis develop, in order to provide the maximum benefit.



Pour the entire contents of the single-dose Gelclair<sup>®</sup> sachet into a glass and add up to 40mL or two tablespoons of water. If water is not available, Gelclair<sup>®</sup> may be used undiluted.



Stir mixture and use at once.



Rinse around mouth for at least one minute or as long as possible to coat tongue, palate, throat, inside of cheeks and all oral tissue thoroughly.



Gargle and spit out – do not swallow, although if accidentally swallowed, no side-effects are anticipated.



Use at least three times per day or as needed.

Do not eat or drink for at least one hour following treatment.

If patients are performing oral care (e.g. brushing teeth), it should be performed prior to using Gelclair<sup>®</sup>.

If being treated with other topical agents (analgesic or anti-infective) they should be used before Gelclair<sup>®</sup> since the barrier it forms may prevent other topical agents from reaching the oral mucosa.

Store below 25°C. Do not refrigerate. Keep out of reach of children.

GELCLAIR<sup>®</sup> is a trademark of Helsinn Healthcare SA, Switzerland.

Manufacturer: Biokosmes Srl, Lecco, Italy.

Gelclair<sup>®</sup> is distributed by Orphan Australia Pty. Ltd.,

48 Kangan Drive Berwick, Victoria 3806 Australia under licence from Helsinn Healthcare SA, Switzerland.

Phone: 03 9769 5744 Fax: 03 9769 5944 Website: [www.orphan.com.au](http://www.orphan.com.au)

OA/GELCLAIR/CARDS/0407 ETAL1963ORP



**HEL SINN**